

EUVE OBSERVATIONS: ATMOSPHERES OF COOL BINARIES

NASA Grant NAG5-2330

Annual Report No. 2

For the period 1 August 1995 through 31 July 1996

Principal Investigator

Dr. Andrea K. Dupree

October 1996

Prepared for

National Aeronautics and Space Administration

Washington, D.C. 20546

Smithsonian Institution
Astrophysical Observatory
Cambridge, Massachusetts 02138

The Smithsonian Astrophysical Observatory
is a member of the
Harvard-Smithsonian Center for Astrophysics

The NASA Technical Officer for this Grant is Dr. Ronald J. Oliverson, Code 684.1, Laboratory for Astronomy and Solar Physics, Space Sciences Directorate, NASA/Goddard Space Flight Center, Greenbelt, MD 20771.

Publications and Presentations

Dissecting the EUV Spectrum of Capella, N. S. Brickhouse, 1996, in IAU Colloquium No. 152, Astrophysics in the Extreme Ultraviolet, S. Bowyer and R. F. Malina (eds), Kluwer:Boston, 105–112.

EUV Spectroscopy of Stellar Coronae, N. S. Brickhouse 1996, to appear in *Proceedings of the Tenth APS Topical Conference on Atomic Processes in Plasmas*.

Atomic Processes in Astrophysics, J. C. Raymond & N. S. Brickhouse 1996, *Astrophys. and Space Science*, **237**, 321.

EUVE & ASCA Spectra of Coronal Sources, N. S. Brickhouse, May 1996, presented at AAS HEAD Conference, San Diego.

Uncertainties in Plasma Emission Models for X-ray Spectroscopy, N. S. Brickhouse, May 1996, presented at AAS HEAD Conference Workshop, Information from X-Ray Spectra, San Diego.

High Temperature Structure in Cool Binary Stars, A. K. Dupree, N. S. Brickhouse, & G. J. Hanson, 1996, IAU Colloquium No. 152, Astrophysics in the Extreme Ultraviolet, S. Bowyer and R. F. Malina (eds), Kluwer:Boston, 141–145.

Coronal Structure in Capella, A. K. Dupree, and N. S. Brickhouse, in IAU Symposium No. 176: Stellar Surface Structure, Poster Proceedings, University of Vienna, 184–186.

The EUV Transition Lines of Capella, N. S. Brickhouse, A. K. Dupree, & J. C. Raymond, in 9th Cambridge Workshop on Cool Stars, Stellar Systems and the Sun, ASP Conf. Series, R. Pallavicini & A. K. Dupree (eds), in press.

EUVE Spectroscopy of Active Binaries, A. K. Dupree, in 9th Cambridge Workshop on Cool Stars, Stellar Systems and the Sun, ASP Conf. Series, R. Pallavicini & A. K. Dupree (eds), in press.

EUVE Observations of a Giant Flare from UX Arietis, A. K. Dupree & N. S. Brickhouse, BAAS, 28, 942, 1996

EUVE Observations of 44i Boo: Coronal Structure and Variability, N. S. Brickhouse & A. K. Dupree, BAAS, 28, 942, 1996.

Dissecting the EUV spectrum of *Capella*

By NANCY S. BRICKHOUSE¹

¹Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

Extreme ultraviolet spectra of *Capella*, obtained at various orbital phases over the past two years by the *EUVE* satellite, show strong emission lines from a continuous distribution of temperatures ($\sim 10^5$ – $10^{7.3}$ K). In addition to the strong He II $\lambda 303.8$, the spectra are dominated by emission lines of highly ionized iron. Strong lines of Fe IX, XV, XVI, and XVIII–XXIV are used to construct emission measure distributions for the individual pointings, which show several striking features, including a minimum near 10^6 K and a local maximum at $10^{6.8}$ K. Furthermore, intensities of the highest temperature lines ($T_e > 10^7$ K) show variations (factors of 2–3) at different orbital phases, while the lower temperature Fe lines show variations of about 30% or less. The low variability of most of the strong low temperature features motivates a detailed analysis of the summed spectrum. With ~ 280 ksec of total exposure time, we have measured over 200 emission features with $S/N \geq 3.0$ in the summed spectrum. We report here initial results from the analysis of this spectrum. We can now identify lines of Fe VIII and X–XIV, as well as a number of electron density and abundance diagnostic lines.

We also report here the first *direct* measurement of the continuum flux around ~ 100 Å in a cool star atmosphere with *EUVE*. The continuum flux can be predicted from the emission measure model based on Fe line emission, and demonstrates that the Fe/H abundance ratio is close to the solar photospheric value.

1. Introduction

Capella (Alpha Aurigae; HD 34029) is a bright nearby multiple star system that has been the target of numerous observations over the past two decades in the UV and X-ray regions. Under the Guest Observer Program of the *EUVE* satellite, we have now obtained multiple observations of this source. All of the individual pointings confirm the initial results of the calibration data analysis, reported by Dupree *et al.* (1993): (a) a continuous distribution of temperatures from the transition region to the hot corona; (b) a minimum in the emission measure distribution (EMD) near 10^6 K; (c) a local maximum in the EMD at $10^{6.8}$ K; and, (d) a number of transition region lines that are weak relative to their ultraviolet counterparts. It is the goal of this work to explore the predictive capability of the EMD derived from strong Fe lines. Using the summed spectrum from all the *EUVE* observations, we have now measured over 200 emission features with signal to noise (S/N) ≥ 3 in the three spectrometers. A more detailed analysis can be found in Brickhouse *et al.* (1995b).

The analysis of strong EUV Fe lines reveals complex structure in the *Capella* atmosphere not previously discerned by low resolution X-ray spectroscopy. For example, Swank *et al.* (1981) characterize a number of cool binary coronae as two-temperature distributions using observations taken with the *Einstein* Solid State Spectrometer. Even the moderate resolution *EXOSAT* Transmission Grating Spectrometer was unable to resolve line blends, and thus the multithermal emission models of Lemen *et al.* (1989) are simple parameterizations reliant on global spectral fitting. Among the early results from *EUVE* spectroscopy are clean, strong emission lines of Fe IX, XV, XVI, and XVIII–XXIV in a number of bright active binaries (Dupree *et al.* 1993; Landini & Monsignori Fossi 1993; Stern *et al.* 1995). For example, Dupree *et al.* (1993) report about twenty Fe lines from which an EMD is constructed. While the *Capella* distribution is indeed reminiscent

of two dominant temperatures (unlike certain other sources), a continuous distribution is required to fit the *EUVE* lines.

Given the sharpness of features in *Capella*'s atmosphere and the rich diversity of structures found in other sources (Dupree *et al.* 1995), a number of questions arise as to the physical conditions at various temperatures in the transition region and corona. Dupree *et al.* (1993) find evidence for high densities (10^{12} to 10^{13} cm $^{-3}$) from N_e -sensitive lines of Fe XXI. Brickhouse *et al.* (1995a) discuss apparent inconsistencies in the densities derived from different line ratios, suggesting that some combination of weak lines, blends, and different source regions for Fe XXI might be responsible. Since these densities at $T_e \sim 10^7$ K imply small emitting volumes and confining magnetic fields of several hundred gauss, the confirmation of such high densities is a strong motivation for longer observations.

At lower temperatures, the EMD has been much less well constrained, relying solely on Fe IX below $T_e \sim 2 \times 10^6$ K to connect the transition region to the corona. Since the emission measure minimum is at a significantly higher temperature ($\sim 10^6$ K) than that of the sun ($\sim 10^5$ K), the role of various components in the energy balance of *Capella* must be quite different from the sun. The summed spectrum now fills in much of the gap in coverage of Fe ionization stages, adding Fe VIII and X–XIV. Many of these lines are also N_e -sensitive.

Drake *et al.* (1995) discuss the importance of elemental abundance measurements in stellar coronae for understanding coronal heating, and as Cook *et al.* (1989) point out, significant variations from solar photospheric abundances can also change the shape of the radiative loss function. The summed spectrum of *Capella* includes lines from many of the abundant elements; we report here the analysis of emission formed at high temperature from elements other than Fe. Discussion of EUV lines formed in the transition region and near the emission measure minimum is forthcoming (Brickhouse *et al.* 1995b).

2. Observations and data reduction

The individual spectra were acquired over five separate pointings. The extraction techniques, described elsewhere (Dupree *et al.* 1995a), have been applied to the individual observations separately. The spectra are co-added to produce the summed spectra for a total of 2.82, 2.77, and 2.83 ksec for the SW, MW, and LW spectrometers, respectively. Figure 1 shows these spectra, with the S/N curve overplotted. The S/N of individual lines is calculated over the entire line profile, and thus is somewhat better than the peak value shown. Since the individual spectra were accumulated under different conditions, the simple summation leads to a minor degradation of the effective spectral resolution. The strong, isolated spectral lines maintain Gaussian profiles, with fits that are generally improved more by higher S/N than they are degraded by decreased spectral resolution; they are consistent with the performance characteristics described by Boyd *et al.* (1994). For individual lines, the centroid and flux determinations, the continuum flux subtraction and the correction for interstellar absorption are done as in Dupree *et al.* (1993). For blended lines, we require that the line widths, deviations from laboratory wavelengths, and wavelength separations of the individual lines be consistent with the spectrometer characteristics before accepting line identification and fluxes.

Variations during the individual pointings are at the 20% level. Dupree *et al.* (1995b) find that the intensities of the highest temperature lines ($T_e > 10^7$ K) show modest variations (factors of 2 to 3) with different pointings, while the lower temperature lines ($T_e < 10^7$ K) show variations of about 30% or less. Ayres (1988) shows that the UV line fluxes are remarkably constant with time as well. While variability in the high




FIGURE 1. *Capella* summed spectrum in the three *EUVE* spectrometers (solid line); signal to noise (dotted line). Curves are smoothed by 5 and 7 pixels, respectively.

temperature lines is useful for line identification and deblending (such as in the blend of Fe XXIV, Ca XVII, Fe XII, and O V around 192Å), the lack of variability over most of the temperature range gives confidence that this analysis is appropriate for typical atmospheric conditions.

3. The EMD for the summed spectrum

The analysis of a coronal source through its EMD provides key insights into heating, cooling, and confinement. While these issues are central to much of the EUV work on stellar coronae, the more immediate goal here is to use the predictive capabilities of an EMD, determined by strong lines from a single element, to examine the rich spectrum for other diagnostics. This goal sets the criterion for the “best fit” as the agreement of strong lines with the model. Figure 2 shows our model for the *Capella* spectrum, and the agreement of the lines used in constructing it.

For $T_e \geq 5 \times 10^5$ K the EMD depends on the the EUV lines of highly ionized Fe. Newly identified lines are discussed by Brickhouse *et al.* (1995b). Although the high temperature extent is not well constrained using EUV lines, this model is now consistent with the *ASCA* continuum above 2 keV (Singh 1994), as well as with the carefully deblended doublet lines of Fe XXIV. We include the X-ray fluxes observed by Vedder & Canizares (1983) in Figure 2, although we do not use these lines in the fitting. It is interesting to note that the two lines formed predominantly at $T_e = 10^{6.8}$ K (O VIII $\lambda 18.97$ and Fe XVII $\lambda 15.01$) are in good agreement with the model, while the higher temperature line (Fe XX $\lambda 12.83$) exceeds the prediction of the summed spectrum, but is consistent within the range of variability.




FIGURE 2. *Capella* model. *Upper*: log Emission Measure (cm^{-3}) vs $\log T_e$ (K). *Lower*: comparison of observed to predicted fluxes for the strong lines. See text for sources of X-ray and UV lines from different elements.

Below this temperature we use the total UV line fluxes measured by Linsky *et al.* (1995) with *HST* and Hurwitz *et al.* (1995) with *ORPHEUS*. As Linsky *et al.* (1995) discuss, the N V resonance line flux may reflect N enrichment in the evolved star, and thus we have forced our fit to the weaker O V line. Many EUV transition region lines are extremely weak relative to the prediction based on the UV emission lines, as initially reported in Dupree *et al.* (1993) and discussed in more detail in Brickhouse *et al.* (1995b).

We cannot emphasize too strongly that the features in the EMD are required by the fluxes of strong lines. For example, *Capella*'s local maximum at $10^{6.8}$ K is determined by eight strong lines of Fe XVIII and XIX, for a total of 7600 counts in the summed spectrum. Continuous distribution models which smooth over this feature will underpredict the emission from other lines formed predominantly at this temperature. On the other hand, a two-temperature model has poor predictive powers for lines formed away from the two dominant temperatures. While the single emission measure at $10^{6.8}$ K (i.e. the integral over the interval $\Delta T_e = \pm 0.05$ dex) in our model accounts for about 80% of Fe XVIII and XIX line fluxes (with most of the rest of the flux coming from $10^{6.7}$ K), it only accounts for 43%, 13%, and 1% of the Fe XVI, XV, and XIV line fluxes, respectively. Thus many temperatures are required to produce the range of Fe ionization stages.

With high S/N spectra and excellent flux calibration, errors in the atomic models begin to dominate the uncertainties. For the more highly ionized Fe species, some collision strengths are as accurate as 10%; for the lower ionization stages (Fe VIII to XIV), collision strengths may be inaccurate at the 30% level or worse. Dielectronic recombination rates are the largest source of uncertainty in the ionization balance, as discussed by Brickhouse *et al.* (1995a). In addition to these uncertainties, many of the lower temperature Fe lines are N_e -sensitive, and may complicate the emission measure analysis. We find from a synthesis of the Fe XII–XIV spectra that $N_e \sim 10^9 \text{ cm}^{-3}$ is more consistent with observations than 10^{10} or 10^{11} cm^{-3} . While we are not able to measure unblended line ratios from these ions, we are able to isolate several line groups.

At high temperature, line ratio diagnostics give $\log N_e$ (cm^{-3}) = 11.1 (+0.3, -no constraint) from Fe XXI $\lambda 128.7/(\lambda 97.9 + \lambda 102.2)$; 12.5 (+0.2, -0.1) from Fe XXI $\lambda 128.7/\lambda 113.3$; 12.35 (+0.2, -0.3) from Fe XXI $\lambda 128.7/\lambda 142.2$; 11.73 (+0.2, -0.3) from Fe XXI $\lambda 128.7/\lambda 145.6$; and, 12.05 (+0.3, -no constraint) from Fe XXII $\lambda 117.1/\lambda 114.4$. Uncertainties given correspond to 1- σ errors in the line ratios. Although the summed spectra provide a great improvement in S/N over the calibration data (Dupree *et al.* 1993), and allow reasonable separation of the blended lines, a large spread remains, which seems to require multiple densities at the source. As noted in Brickhouse *et al.* (1995a), Fe XXI in the model is formed about equally at $T_e = 10^{6.8}$ K and at $T_e > 10^7$ K.

4. Measurement of the continuum and the Fe/H abundance ratio

Stern *et al.* (1995) measure the continuum emission for *Algol* by summing the flux in 4 bins to achieve sufficient S/N. We are able to measure *directly* the minimum flux level in the *Capella* summed spectrum, since $S/N > 3.0$ in every bin throughout the wavelength range 81-123 Å. Figure 3 is a detail of this spectral region. We attribute the minimum flux to real continuum emission (mostly bremsstrahlung), and compare it to model predictions to derive an Fe/H abundance ratio. We find that the Fe/H abundance ratio is 0.88 ± 0.13 solar, assuming the solar values of Anders & Grevesse (1989).

The flux in the pixels which contain the lowest flux values determines the continuum, and the quandary is to select the appropriate number of pixels in the flux averaging. We select the lowest points in the spectrum that agree in value with each other to within their statistical error limits. With smoothing by 3 pixels, this condition produces 14 bins to define the continuum, and the derived error is the standard deviation. The value of the continuum defined in this way is consistent with the measurement determined (from fewer points) with smoothing by 5 or 7 pixels. Furthermore, inspection of line lists gives additional confidence that emission lines are avoided. For all but 3 bins, there are no close wavelength matches with lines listed in the Doschek & Cowan (1984) line list, which includes lines from flaring as well as other solar conditions. We can estimate the contribution from weak, highly ionized Ni by scaling from lines we observe relative to their theoretical Fe counterparts, for which the tokamak line lists of Davé *et al.* (1987) and Stratton *et al.* (1985) are invaluable. The existence of pseudo-continua produced by unknown weak lines is difficult to rule out without models for these weak lines, but their contribution should be well within the errors on the continuum flux measurement.

Figure 3 also shows the temperature-dependence of bremsstrahlung emission. In our model the emission comes predominantly from the $10^{6.8}$ K local maximum. The model predicts that the continuum emission decreases by about 20% over the range from 85 to 140 Å; we are not able to determine a slope observationally.

5. Elemental abundances at high temperature

The lines used to determine abundances are listed in Table 1. Line emissivities are from Raymond (1988), modified by the atomic data of Zhang *et al.* (1990) for Li-like ions. Figure 4 shows the dependences of model line emissivities and integrated intensities. For the high Z (above Ne) Li-like lines, the emission in our model comes primarily from $10^{6.8}$ K. The doublet lines are particularly useful, since their line ratio can give additional confidence in the flux measurement. The O and Ni lines in the Table similarly come from this temperature in the model. The errors listed are observational, but there are systematic errors as well, particularly errors in the correction for interstellar absorption. The effect of assuming neutral He is comparable to the errors in N_H given




FIGURE 3. *Left*: continuum emission in flux units, where 1 pix = 0.067 Å. Dotted vertical lines indicate the wavelength range for which $S/N > 3$. *Right*: model of continuum flux at 100 Å (0.12 keV) (solid) and at 6 Å (2.0 keV) (dash-dotted) vs temperature. The contribution fraction is the integral of the continuum emissivity over the EMD relative to the total.




FIGURE 4. The fractional emissivity (dashed) and intensity (solid) contributions as functions of temperature for selected lines from Li-like ions.

by Linsky *et al.* (1993). Vennes *et al.* (1993) show that the fraction of ionized He might be as high as 25%. The abundances in Table 1 derived from lines between 227 and 504 Å would then be somewhat reduced. For Si XII the relative abundance would decrease from 1.99 to 1.65, and would bring the doublet ratio into better agreement with theory.

6. Conclusions

High quality *EUVE* spectra of cool stars, such as the summed spectrum of *Capella*, are rich with information on the physical conditions in the atmosphere. Long observations confirm the presence in *Capella* of high densities ($\geq 10^{12} \text{ cm}^{-3}$) at high temperature, but lower densities are present as well at other temperatures. We have now measured the relative abundances of a number of elements which emit radiation at the local maximum

Element	Emission Lines	<i>Capella</i> [†]	<i>Capella</i> [‡]	Solar Corona [¶]
O	O VIII $\lambda 102.45^{\parallel}$	0.42 ± 0.18	0.13(0.03,0.27)	0.9
	O VIII $\lambda 18.97^{\dagger\dagger}$	1.02 ± 0.28		
Si	Si XII $\lambda 499.40$	1.99 ± 0.38	0.85(0.67,1.07)	3.6
	Si XII $\lambda 520.67$			
S	S XIV $\lambda 417.61$	1.15 ± 0.18	0.73(0.39,1.14)	1.1
	S XIV $\lambda 445.77$			
Ar	Ar XVI $\lambda 353.92$	2.41 ± 0.31		1.0
	Ar XVI $\lambda 389.14$			
Fe	(all)	0.88 ± 0.13	0.46(0.36,0.61)	2.7
Ni	Ni XVII $\lambda 249.18$	1.81 ± 0.20		3.9
	Ni XVIII $\lambda 291.97$			
	Ni XVIII $\lambda 320.54$			

TABLE 1. Elemental Abundances at $T_e \sim 10^{6.8}$ K^{‡‡}

[†] This work. Errors are 1- σ estimates of measurement uncertainties, including statistical errors, deblending of weak lines and high order flux subtraction.

[‡] From Drake *et al.* (1994). Allowed ranges reflect 90% confidence levels from X-ray spectral fitting with two temperatures.

[¶] Feldman (1992)

^{||} This line was deblended from Fe XXI $\lambda 102.2$.

^{††} Observed by Vedder & Canizares (1983). Prediction uses the EMD from this work.

^{‡‡} Abundances are relative to solar photospheric abundances of Anders & Grevesse (1989).

at $\sim 10^{6.8}$ K, and find that they are more consistent with solar photospheric than coronal values. Density and abundance measurements may provide keys to understanding the nature of this enhanced emission region. Variability in the N_e -sensitive lines at the highest temperatures may provide a tool for distinguishing different sources of emission.

The author appreciates the contributions of Andrea Dupree, John Raymond and Greg Hanson. This work has been supported by NAGW-528 and NAG5-2330 to the Smithsonian Institution.

REFERENCES

- ANDERS, E., & GREVESSE, N. 1989 Abundances of the Elements: Meteoritic and Solar. *Geochim. Cosmochim. Acta* **53**, 197–214.
- AYRES, T. R. 1988 A Spectral Dissection of the Ultraviolet Emissions of *Capella*. *Ap. J.* **331**, 467–476.
- BOYD, W., JELINSKY, P., FINLEY, D. S., DUPUIS, J., ABBOTT, M., CHRISTIAN, C., & MALINA, R. F. 1994 In-Orbit Performance of the Spectrometers of the Extreme Ultraviolet Explorer. *Proc. SPIE*, **2280**, 280–296.
- BRICKHOUSE, N. S., RAYMOND, J. C., & SMITH, B. W. 1995a New Model of Iron Spectra in the Extreme Ultraviolet and Application to *SERTS* and *EUVE* Observations: A Solar Active Region and *Capella*. *Ap. J. Supp.* **97**, 551–570.
- BRICKHOUSE, N. S., DUPREE, A. K., & RAYMOND, J. C. 1995b Analysis of the *EUVE* Spectrum of *Capella* from the Transition Region to the Hot Corona. In progress.

- COOK, J. W., CHENG, C.-C., JACOBS, V. L., & ANTIOCHOS, S. K. 1989 Effect of Coronal Elemental Abundances on the Radiative Loss Function. *Ap. J.* **338**, 1176–1183.
- DAVÉ *et al.* 1987 Time-resolved spectra in the 80–340 Å Wavelength Region from Princeton Large Torus Tokamak Plasmas., *J. Opt. Soc. Am.* **B4**, 635–643.
- DOSCHEK, G. A., & COWAN, R. D. 1984 A Solar Spectral Line List Between 10 and 200 Å Modified for Application to High Spectral Resolution X-ray Astronomy. *Ap. J. Supp.* **56** 67–89.
- DRAKE, J. J., LAMING, J. M., & WIDING, K. G. 1995 Stellar Coronal Abundances. II. The First Ionization Potential Effect and Its Absence in the Corona of Procyon. *Ap. J.* **443**, 393–415.
- DRAKE, S. A., SINGH, K. P., WHITE, N. E., & SIMON, T. 1994 *ASCA* X-ray Spectra of the Active Single Stars β Ceti and Π^1 Ursae Majoris. *Ap. J.* **436**, L87–L90.
- DUPREE, A. K., BRICKHOUSE, N. S., DOSCHEK, G. A., GREEN, J. C., & RAYMOND, J. C. 1993 The Extreme Ultraviolet Spectrum of Alpha Aurigae (*Capella*). *Ap. J.* **418**, L41–L44.
- DUPREE, A. K., BRICKHOUSE, N. S., & HANSON, G. J. 1995a High Temperature Structure in Cool Binary Stars. This proceedings.
- DUPREE, A. K., BRICKHOUSE, N. S., DOSCHEK, G. A., HANSON, G. J., & RAYMOND, J. C. 1995b EUVE Spectra of Alpha Aurigae (*Capella*) at Different Phases. In preparation.
- FELDMAN, U. 1992 Elemental Abundances in the Upper Solar Atmosphere. *Phys. Scripta* **46**, 202–220.
- HURWITZ, M., BOWYER, S., & DUPREE, A. K. 1995 *ORPHEUS* Observations of *Capella*. In preparation.
- LANDINI, M., & MONSIGNORI FOSSI, B. C. 1993 Extreme Ultra Violet Plasma Diagnostic: a Test Using EUVE Calibration Data. *Astron. Astrophys.* **275**, L17–L20.
- LEMEN, J. R., MEWE, R., SCHREIVER, C. J., & FLUDRA, A. 1989 Coronal Activity in F-, G-, and K-Type Stars. III. The Coronal Differential Emission Measure Distribution of *Capella*, σ^2 CrB, and Procyon. *Ap. J.* **341**, 471–483.
- LINSKY, J. L. *et al.* 1993 Goddard High-Resolution Spectrograph Observations of the Local Interstellar Medium and the Deuterium/Hydrogen Ratio along the Line of Sight toward *Capella*. *Ap. J.* **402**, 694–709.
- LINSKY, J. L., WOOD, B. E., JUDGE, P., BROWN, A., ANDRULIS, C., & AYRES, T. R. 1995 The Transition Regions of *Capella*. *Ap. J.*, **442**, 381–400.
- RAYMOND, J. C. 1988 *Radiation from Hot, Thin Plasmas*. in R. Pallavicini, ed, *Hot Thin Plasmas in Astrophysics* (Dordrecht: Kluwer), 3–20.
- SINGH, KP 1994, private communication.
- STERN, R. A., LEMEN, J. R., SCHMITT, J. H. M. M., & PYE, J. P. 1995 EUVE Observations of Algol: Detection of a Continuum and Implications for the Coronal [Fe/H] Abundance. *Ap. J.*, **444**, L45–L48.
- STRATTON *et al.* 1985 Relative Intensities of $2s^2\ 2p^k$ – $2s\ 2p^{k+1}$ transitions in FI- to BI-like Ti, Cr, Fe, Ni, and Ge in a Tokamak Plasma: A Comparison of Experiment and Theory., *Phys. Rev. A* **31**, 2534–2547.
- SWANK, J. H., WHITE, N. E., HOLT, S. S., & BECKER, R. H. 1981 Two-Component X-ray Emission from RS Canum Venaticorum Binaries. *Ap. J.* **246**, 208–214.
- VEDDER, P. W., & CANIZARES, C. R. 1983 Measurement of Coronal X-ray Emission Lines from *Capella*. *Ap. J.* **270**, 666–670.
- VENNES, S. *et al.* 1993 The First Detection of Ionized Helium in the Local ISM: *EUVE* and *IUE* Spectroscopy of the Hot DA White Dwarf GD 246. *Ap. J.* **410**, L119–L122.
- ZHANG, H. L., SAMPSON, D. H., & FONTES, C. J. 1990 Relativistic Distorted-Wave Collision Strengths and Oscillator Strengths for the 85 Li-like Ions with $8 \leq Z \leq 92$. *At. Data Nucl. Data Tables*, **44**, 31–77.

High Temperature Structure in Cool Binary Stars

By A. K. Dupree¹, N. S. Brickhouse¹
AND G. J. Hanson¹

¹Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138 USA

Strong high temperature emission lines in the EUVE spectra of binary stars containing cool components [Alpha Aur (*Capella*), 44 ϵ Boo, Lambda And, and VY Ari] provide the basis to define reliably the differential emission measure of hot plasma. The emission measure distributions for the short-period ($P \leq 13$ d) binary systems show a high temperature enhancement over a relatively narrow temperature region similar to that originally found in *Capella* (Dupree *et al.* 1993). The emission measure distributions of rapidly rotating single stars 31 Com and AB Dor also contain a local enhancement of the emission measure although at different temperatures and width from *Capella*, suggesting that the enhancement in these objects may be characteristic of rapid rotation of a stellar corona. This feature might be identified with a (polar) active region, although its density and absolute size are unknown; in the binaries *Capella* and VY Ari, the feature is narrow and it may arise from an interaction region between the components.

1. Introduction

The emission measure defined by the well-exposed EUV spectra of the binary system *Capella* (G8 III + G0 III) surprisingly revealed a *narrow* high temperature “bump” (Dupree *et al.* 1993) occurring at $T = 6.3 \times 10^6$ K. The feature is continuously present irrespective of the orbital phase of the system (Dupree *et al.* 1994; 1995), and may arise from a continuously visible hot region on one component, or from an interacting region between the stars. The distribution of the high temperature emission measure in the *Capella* system differs from the broad emission measure distribution found in a solar active region (see Brickhouse *et al.* 1995). We investigate whether this feature is characteristic of binary systems and whether rotation controls its presence.

2. Spectra and the Emission Measure Distribution

EUVE spectra from our Guest Observer program and from the EUVE Science Data Archives were obtained for 4 binary systems (VY Ari, *Capella*, 44 ϵ Boo, λ And) and two rapidly rotating single stars (AB Dor and 31 Com). EUV spectra were extracted from the images and calibrated in a uniform manner. The reduced spectra are shown in Fig. 1. In spite of the varying physical characteristics of the stars, there are similarities among the spectra. The resonance line of He II ($\lambda 304$) dominates the MW spectra; strong resonance lines of iron dominate the SW spectra. However, the relative strengths of the lines differ among these objects, and so require detailed evaluation of the emission measure. It is the strength of the Fe XVIII and Fe XIX transitions relative to Fe XX, XXIII that controls the presence of the locally enhanced emission measure (the “bump”) in *Capella*.

Signal to noise ratios were evaluated at the EUVE spectral resolution, and only the strongest features ($S/N \approx 9 - 15$) were selected to determine the emission measure. Correction of the observed fluxes for interstellar absorption was made with N_H from Table 1. The H/He abundance ratio was set to 11.6 (Kimble *et al.* 1993) and helium was assumed to be neutral. Assumption of higher values for N_H would strengthen the




FIGURE 1. EUVE spectra in the SW (*left panel*) and MW (*right panel*) band for the rapidly rotating single stars AB Dor and 31 Com and the binaries 44 Boo, *Capella*, VY Ari, and λ And. Species of Fe are marked by ionization stage. Fe XXIII ($\lambda 132.8$) is blended with Fe XX.

HD	Star	Sp. Type/Lum.	P_{orbital} (days)	$P_{\text{phtn.}}$ (days)	N_H (cm^{-2})	SW Exp (s)
17433	VY Ari†	K3-4V-IV	13.208	16.64	1.0×10^{18}	168557
34029	α Aur‡	G0 III/G8 III	104.0	8/80	1.8×10^{18}	281571
36705	AB Dor¶	K1 IIIp	...	0.514	1.0×10^{18}	166460
111812	31 Com	G0 III	< 6	...	1.0×10^{18}	84503
133640	44 ι Boo††	G0 V	0.268	...	1.0×10^{18}	104145
222107	λ And†	G8IV-III	20.52	53.95	4.0×10^{18}	105701

TABLE 1. PARAMETERS OF TARGET STARS

† RS CVn-type binary.

‡ RS CVn-type binary; EUVE spectra of *Capella* taken at several phases were averaged together.

¶ Rapidly rotating single star.

|| 31 Com is a rapidly rotating single star thought to be a Hertzsprung Gap giant. The “orbital” period was evaluated from the $v \sin i$ value of 77 km s^{-1} , assuming a radius of $9R_{\odot}$.

†† W UMa-type contact binary.

Fe XVI emission ($\lambda 336$) relative to the Fe XV emission ($\lambda 284$) were observed, but leave the fluxes in the short wavelength region ($\lambda 85\text{--}\lambda 170$) essentially unchanged.

The emission measure distribution is determined iteratively, by fully integrating the line emissivities through a trial emission measure distribution, and comparing the predicted and observed line fluxes. There is no restriction on the “smoothness” of the emission measure distribution such as occurs with spectral fitting procedures; moreover the fit is not degraded by bins containing noise or weak signals. Only strong lines are considered in evaluating the emission measure. We require that these fluxes agree within acceptable uncertainties which in this case is better than a factor of two. Detailed current models for all iron ions (Brickhouse *et al.* 1995) were used. Results for λ And are taken from another publication (Hanson *et al.* 1995) where fluxes of ultraviolet lines measured with *IUE* were included so that the emission measure below $2 \times 10^5 \text{ K}$ is well-defined.

3. Conclusions

All of the target stars exhibit a range of high ion stages indicating a generally continuous distribution of temperature. The spectral resolution of EUVE demonstrates that two-temperature fits which rely on energy band measures do not represent these stellar atmospheres. For the weaker spectra, the strongest observable lines arise from high stages of ionization (Fe XVIII ... XXIII), the Fe XV ($\lambda 284$) and Fe XVI ($\lambda 336$ and $\lambda 361$) emission is not prominent, and the emission measure can only be reliably defined in a restricted temperature range.

Inspection of the emission measure distributions contained in Fig. 2 and 3 shows that the *Capella* “bump” is not unique. The binary systems (VY Ari, 44 ι Boo) clearly show the enhanced emission measure distribution similar to *Capella*. The appearance of the emission measure enhancement is most pronounced in binary systems with short orbital period (≤ 13 days) and/or high rotational velocity. The orbital period of *Capella* is 104 days, however the hot component of *Capella* is rotating rapidly with a period of ≈ 8 days (Fekel *et al.* 1986) so that the system may be classified as a rapid rotator for these purposes. The strong lines of Fe XVIII ($\lambda 93.9$) and Fe XIX ($\lambda 108.4$) demand the presence





FIGURE 2. Emission measure distribution determined for 4 binary systems.

FIGURE 3. Emission measure distribution determined by the strongest lines for 2 rapidly rotating single stars (31 Com and AB Dor) and the binary system *Capella*.

of an increased emission measure in *Capella* for their production. That *44 Boo* would have a similar feature is apparent from inspection of its SW spectrum; the enhancement is broader in temperature than the *Capella* bump. The SW spectrum of *VY Ari* shows an exceptionally strong feature of Fe XX, XXIII ($\lambda 132.8$) relative to the Fe XVIII and XIX transitions; thus the local enhancement of the emission measure occurs at a temperature higher by 0.2 dex, namely $T(K) = 7.0$ dex. The long period system, λ And has a much

less well defined emission measure enhancement (Hanson *et al.* 1995). The slopes of the emission measures with temperature are consistent with models of magnetic loops or arcades.

Two single stars can address the question of whether the “bump” is unique to binaries. The rapidly rotating single giant 31 Com is of particular interest because it lies in the Hertzsprung gap (Wallerstein *et al.* 1994). Since the hot rapidly rotating secondary in the *Capella* system is believed (Pilachowski & Sowell 1992) to be passing through the Hertzsprung gap too, 31 Com might prove helpful in disentangling the composite EUV spectrum of *Capella*. The star 31 Com has a broad enhancement required by the dominant Fe XXII and Fe XXIII transitions, at a temperature higher (7.0 dex) than the *Capella* feature. Such a broad hot component is not found in the *Capella* emission measure. AB Dor, another rapidly-rotating single giant shows a broad enhancement, also different from *Capella*. Thus, rapidly rotating single stars have “bumps” but they appear different from those in *Capella* and VY Ari.

Rotation appears to be a significant physical parameter in producing an enhanced emission measure feature in cool star atmospheres. This feature is generally present in those systems with periods ≤ 13 days. However the temperature of the maximum and the width and strength of the enhancement differ from star to star. It is plausible to associate this feature with magnetic structures in a rapidly-rotating corona, in which case it could be dense and of small scale; however, in the binary systems *Capella* and VY Ari, a narrow enhancement (of unknown density) is found. It may be that such a feature results from interaction between the components. Binaries of longer period, represented by λ And and σ Gem (Hanson *et al.* 1995) do not show such well-defined enhancements in the emission measure, although they possess equally hot plasma. However longer EUV exposures are needed on many of these stars to detect weaker species and to define the atmospheric structure over a wider temperature range.

This work is partly supported by NASA NAG5-2330 to the Smithsonian Institution.

REFERENCES

- BRICKHOUSE, N. S., RAYMOND, J. C., & SMITH, B. W. 1995 *New Model of Iron Spectra in the Extreme Ultraviolet and Application to SERTS and EUVE Observations of a Solar Active Region and Capella*. *Ap. J. Suppl.*, **97**, 551-570.
- DUPREE, A. K., BRICKHOUSE, N. S., DOSCHEK, G. A., GREEN, J. C., & RAYMOND, J. C. 1993 *The Extreme Ultraviolet Spectrum of Alpha Aurigae (Capella)*. *Ap. J.*, **418**, L41-L44.
- DUPREE, A. K., BRICKHOUSE, N. S., DOSCHEK, G. A., HANSON, G. J., & RAYMOND, J. C. 1994 *EUVE Spectra of Alpha Aurigae (Capella) at Different Phases*. *B.A.A.S.*, **26**, 864.
- DUPREE, A. K., BRICKHOUSE, N. S., & HANSON, G. J. 1995, *Inhomogeneity of Coronal Structures in Capella*, in preparation.
- FEKEL, F. C., MOFFETT, T. J., & HENRY, G. W. 1986 *A Survey of Chromospherically Active Stars* *Ap. J. Suppl.*, **60**, 551-576.
- HANSON, G. J., BRICKHOUSE, N. S., & DUPREE, A. K. 1995 *EUVE Spectra of Lambda Andromedae and Sigma Geminorum*. *Ap. J.*, submitted.
- KIMBLE, R. PLUS 13 AUTHORS 1993 *Extreme Ultraviolet Observations of G191-B2B and the Local Interstellar Medium with the Hopkins Ultraviolet Telescope*. *Ap. J.*, **404**, 663-672.
- PILACHOWSKI, C. A., & SOWELL, J. R. 1992 *The Lithium Abundance of the Capella Giants*. *A. J.*, **103**, 1668-1672.
- WALLERSTEIN, G., BOHM-VITENSE, E., VANTURE, A. D., & GONZALEZ, G. 1994 *The Lithium Content and Other Properties of F2-G5 Giants in the Hertzsprung Gap*. *A. J.*, **107**, 2211-2221.